



[Set home page](#) · [Bookmark site](#) · [Add search](#)

[Latest News](#)

[Browse Topics](#)

[Encyclopedia](#)

[Health Center](#)

[Videos](#)

[Amazon Store](#)

Show/Hide Menu

[Print this page](#) | [Email to friend](#)

Text Size > A A A

Front Page

- > Breaking News
- > Today's Digest
- > Week in Review
- > Email Updates
- > **RSS** Newsfeed

News Sections


- > Health & Medicine
- > Mind & Brain
- > Plants & Animals
- > Space & Time
- > Earth & Climate
- > Matter & Energy
- > Computers & Math
- > Fossils & Ruins

Science Topics

- > Agriculture
- > Astronomy
- > Biology
- > Chemistry
- > Earth Sciences
- > Environment
- > Mathematics
- > Physics
- > Social Sciences
- > Technology
- > **more topics**

Health Topics

- > Aging
- > Diseases
- > Fitness



Source: [University of California - Berkeley](#)

Date: 2005-11-10

[Print this page](#)


[Email to friend](#)

New System For Earthquake Early Warning

A University of California, Berkeley, seismologist has discovered a way to provide seconds to tens of seconds of advance warning about impending ground shaking from an earthquake.

While a few seconds may not sound like much, it is enough time for [school](#) children to dive under their desks, gas and electric companies to shut down or isolate their systems, phone companies to reroute traffic, airports to halt takeoffs and landings, and emergency providers to pinpoint probable trouble areas. Such actions can save lives and money.

An early warning system like this is possible thanks to the work of Richard Allen, UC Berkeley assistant professor of earth and planetary science, who in the last five years has demonstrated that within a few seconds of an earthquake rupture, he can predict the total magnitude of the quake and its destructive potential. In San Francisco, for example, Allen estimates that it's likely the city could receive 20 seconds' warning of an impending temblor.



This view along the fault plane of the 1999 Chi-Chi earthquake in Taiwan shows that the 7.6-magnitude rupture started at the star and radiated outward, with

Search Our A

Find: _____
 in: All News S
 from: 1995
 sort: relevant
 > /

New Job Post

Find: _____
 City: _____
 State: ALL
 View: All Jobs by
 Post: [Jobs / Res](#)

- > Medicine
- > Men's Health
- > Mental Health
- > Nutrition
- > Reproduction
- > Senses
- > Women's Health
- > **more topics**

Computing

- > Artificial Intell.
- > Communications
- > Computer Science
- > Graphics
- > Human Interface
- > Internet
- > Robotics
- > Security
- > Supercomputing
- > Virtual Reality
- > **more topics**

Encyclopedia

- > Agriculture
- > Anthropology
- > Archaeology
- > Astronomy
- > Biology
- > Chemistry
- > Communication
- > Computing
- > Earth Science
- > Engineering
- > Health Science
- > Mathematics
- > Physics
- > Psychology
- > Technology
- > **science topics**
- > **medical topics**

Science Shop

- Books ...
- > Science
- > Mind & Body
- > Engineering

"We can determine the magnitude within a couple of seconds of initiation of rupture and predict the ground motion from seconds to tens of seconds before it's felt," Allen said. He and his colleagues are now testing a system, ElarmS, that would make these predictions, and the researchers are working with the U.S. Geological Survey (USGS) to determine how accurate these warnings would be.

Allen and coauthor Erik L. Olson, a former graduate student at the University of Wisconsin, Madison, published their data on early earthquake ground motion predictions in the Nov. 10 issue of Nature.

Seismologists, especially those in the United States, have become increasingly pessimistic about being able to predict earthquakes. Experiments at the intensively monitored Parkfield, Calif., site have dampened enthusiasm that earthquake ruptures could be predicted hours or days before they happen. To reduce loss of life and property, earthquake-prone regions generally rely on a combination of advance preparation and post-earthquake assessment and notification between five and 10 minutes after a quake.

Allen's early warnings come after a quake rupture has already begun but before the shaking is felt tens of miles from the epicenter.

San Francisco, for example, sits about midway along the northern half of the 800-mile San Andreas fault. If a rupture occurs at the extreme northern end, it could take 80 seconds, traveling nearly 2 miles per second, to reach the city. An early warning system could provide a critical buffer for residents, businesses and emergency responders, even if the time isn't sufficient to evacuate a building.

The early warning information also would feed directly into the new active-response building designs that change the mechanical properties of a structure to let it ride out shaking and minimize damage both inside and out. Active response buildings are already operational in Japan, Allen

each circle representing 3 seconds of time. The green, yellow and red colors indicate regions of high slip. Using ElarmS, it was possible to estimate the final magnitude of the earthquake a mere two seconds after it started, when only the area within the red circle had ruptured. (Douglas Dreger/UC Berkeley [2001])

Related News Stories

Earthquake Damages Neighboring Fault, UCLA And USC Scientists Report (February 3, 2003) -- The 1999 magnitude 7.1 Hector Mine earthquake, in the Mojave Desert near Los Angeles, damaged the fault that broke in the 7.3 magnitude Landers earthquake seven years earlier, report John Vidale, ... > [full story](#)

It's Not My Fault: New Study Shakes Up Interest In Earthquakes (November 3, 1998) -- A new look at the information on a 1971 California earthquake shows that several nearby faults were activated during the temblor, supporting evidence from more recent earthquakes that nearby faults ... > [full story](#)

Earthquakes Kill Thousands In 2003; Deadliest Year Since 1990 (January 26, 2004) -- According to the U.S. Geological Survey (USGS), 2003 closed as the deadliest year for earthquakes since 1990, 25 times more fatal than 2002; 43,819 deaths have been reported for the past year, as ... > [full story](#)

New Map Of Daily Probability Of Earthquake Shaking In California (June 21, 2005) -- The U.S. Geological Survey (USGS) has announced the release of new public web pages that show the probability of earthquake shaking in the next 24 hours in California. These maps graphically ... > [full story](#)

> more [related stories](#)

Related section: [Earth & Climate](#)



- > Computers etc.
- > Outdoors & Nature
- > Prof'l & Technical
- > Reference
- Magazines ...**
- > Science & Nature
- > Health & Fitness
- > Engineering
- > Computers etc.
- > Electronics etc.

More ...

- > Electronics
- > Computers
- > Video Games
- > Outdoor Living
- > Camera & Photo
- > Tools & Hardware
- > Toys & Games

About This Site

- > Editorial Staff
- > Awards & Reviews
- > Contribute News
- > Advertise With Us
- > Privacy Policy
- > Terms of Use

said.

"That is our long-term goal, to have the building feel the earthquake, not the occupants," Allen said.

Two years ago, while at the University of Wisconsin, Allen reported differences in the frequency of seismic signals emanating from small and medium earthquakes during the first four seconds of the rupture, with the larger quakes showing lower frequency signals than the smaller quakes. The signal is part of the primary wave, or P wave, that is the first, though least destructive, wave to arrive after a rupture. Most people experience the P wave, which is a pressure wave that travels through rock like sound through air, as a jolt.

This P wave is followed by a secondary wave, or S wave, that shears the ground back and forth and up and down. Shortly after, more destructive surface waves arrive that jerk the ground sideways and later roll in like ocean waves.

In the current [study](#), Allen shows that the relationship between P wave frequency and the total magnitude of the quake holds for major quakes, up to magnitude 8 and higher, as well as for medium and small quakes. Based on the correlation, he can predict the total magnitude of the quake to within 1 magnitude, and for a specific area, like the San Andreas Fault, to within half a magnitude. Magnitude is a measure of the total area that ruptures underground and the average amount of slip along the rupture. A half a magnitude amounts to a factor of 3 difference in ground motion.

"Most seismologists are surprised, and frequently skeptical, that you can predict the magnitude of an earthquake before it has ended, but this is telling us that there is something very different from what we thought about the physics of the processes involved in a rupture," Allen said.

Allen's findings conflict with the current model of earthquake rupture. The "cascade" model assumes that earthquake faults are made up of lots of different-sized patches, each under some degree of stress. When one of the patches is stressed enough to slip, the slip propagates to adjacent patches, which rupture in turn like falling dominoes. The rupture stops only when the stress propagating along the fault zone reaches a patch that is too solidly locked to slip.

Inherent in this model is the idea that the initiating rupture is the same for big and small quakes. Allen's findings suggest this is wrong. Instead, the rupture is different for large and small quakes from the beginning, and the initial rupture contains information that can be used to predict the final size.

He proposes that if the initial rupture generates a large "slip pulse" that travels continuously in all directions



Fanuc for Less
Fanuc Parts in
[Fanucworld.com](http://www.fanucworld.com)

DVT Risks and Symptoms

You or a loved one may be at risk for a condition related to blood clots. Our free risk assessment can help you learn more about the symptoms and treatments for this condition. www.dvt.net

Ez Math

Math Made Easy
Incredible tricks to make math easier and fun! 11yr old girl's inspiration for Unique incredible video - first of its kind in the world of educational videos! Sale: \$39.95 www.ezmatht.com

across the fault plane, the pulse can supply the necessary energy to propagate through patches that would not otherwise have ruptured. Only when the energy in the pulse drops to a level insufficient to overcome the grip of rock on rock does the rupture stop.

"If the rupture pulse initiates with a large slip, it is more likely to evolve into a large earthquake," he and Olson wrote in their report.

Allen's demonstration that this observation holds in earthquakes around the world, from California to Taiwan and Japan, provides a solid basis for constructing an early warning system. Once the magnitude of the quake has been estimated, computers can predict areas of serious ground shaking based on an understanding of a particular fault. Within five seconds, warnings could be sent to cities in the areas calculated to expect damaging ground motion.

Because humans couldn't respond fast enough, Allen said, these warnings would have to rely on computers programmed to respond to quakes of a certain magnitude.

"This allows people to get information about an event before the ground starts shaking and the system goes down," he said.

The ElarmS system also could warn rescue and clean-up personnel of aftershocks, which can cause collapse of unstable debris.

As the rupture proceeds, Allen said, analysis of seismic waves can refine magnitude and ground motion estimates, finally merging into the standard shake map typically produced within minutes of the end of an earthquake.

"We're at the stage where we need to test the accuracy of the system, which we're now doing," Allen said. "Next, we will determine whether the telemetry is fast enough to get data to us within seconds of a rupture."

###

The work was supported by the USGS, University of Wisconsin, Madison, and UC Berkeley.

Editor's Note: The original news release can be found [here](#).

[Apple iPod
White \(16GB\)
MA005LL/A](#)
[Play!](#)



4 GB (Built-in)
1.5 oz., Audio
MP3, WAV, AAC
VBR, AIFF. Dis
1.5 inch

Best Deals

**Best Deal
Name B**



Featured

Free Shipping,
Tax at Amazon

Compare Price
[Amazon](#)

[Beach
Camera.com](#)
[B&H Photo-
Video-Pro](#)

This story has been adapted from a news release issued by University of California - Berkeley.

Ads by Google

Disaster Assistance Loan

Business loan up to \$150,000. Your business must accept VC/MC.

www.Business-Cash-Advance.com

Tsunami Relief Bracelets

Help raise funds for tsunami victims with custom silicone bands.

deluxebracelets.com/tsunami-relief

Help Hurricane Victims

Christian charity helping victims of Hurricane Katrina

www.crossinternational.org

Give to Quake Victims

You Can Help Save Lives Across Asia World Vision - Giving Since 1950

www.WorldVision.org

**We want
to hear from
you!**
Take our
readership

Visit **NewsDaily**
for the latest headlines
around the

|  | Product | Description | Best Deals | Reviews | Search |
|---|---------|-------------|------------|---------|--------|
| Apple iPod nano White (4 GB, MA005LL/A) MP3 Player | | | | | |

Can't find it? Try searching ScienceDaily or the entire web with:



Web sciencedaily.com

Copyright © 1995-2005 ScienceDaily LLC | Contact: editor@sciencedaily.com