

Probabilistic Volcanic Hazard Assessment

Chuck Connor

School of Geosciences
University of South Florida

CIDER, June 2019

PVHA

Background

Conceptual model

Rates

Location


Future

Right: Tephra fallout and kidney disease in Nicaragua. Data courtesy of Kristy Murphy (Texas Children's Hospital)



Top: Panabaj (Guatemala) debris flow (from Charbonnier et al., 2018).

Right: Aso volcano and the Ikata NPP



Japan Court Orders Shutdown of Nuclear Reactor Near Volcano

Dec 14, 2017



Howel Williams (Williams and McBirney, 1979):

- Long term volcanic hazard assessment - primarily based on the geologic record and analogous volcanoes (should take place well in advance of unrest!)
- Short term volcanic hazards assessment - incorporates data on volcanic unrest, uses geophysical signals and related data to forecast the timing and nature of volcanic eruptions.

volcanic hazards in Iceland, based on past events

PVHA


Background

Conceptual
model

Rates

Location

Future



from Gudmundsson et al., Volcanic hazards in Iceland, 2008

PVHA

Background


Conceptual
model

Rates

Location


Future

Conditional probability of inundation by PDCs, Campi Flegrei



from Neri et al., 2015

Are hazards “high” or “low”?



It depends who you are and what your problem is! For example, 10^{-4} annual probability of lahar inundation is a very low hazard for an 80yr old person, and a very high hazard for a 10yr old person.

Cornell (1968), Cornell and Hanks (1994), Stirling et al., (2009)

PVHA


Background

Conceptual model

Rates

Location

Future



$$P[G > g|\Delta t] = P[x, y|\Delta t]P[M|x, y]P[g|M]$$

International Atomic Energy Agency guidelines for site-specific long-term hazard assessment

PVHA


Background

Conceptual model

Rates

Location

Future



see: IAEA (2012) Volcanic Hazards in Site Evaluation for Nuclear Installations. International Atomic Energy Agency, Vienna. IAEA Safety Standards Series No. SSG-21. IAEA (2016) Volcanic Hazard Assessments for Nuclear Installations: Methods and Examples in Site Evaluation. IAEA Techdoc Series No. 1795.

Initial Scoping: Basin and Range volcanism near Yucca Mountain (NV)

PVHA

Background

Conceptual model


Rates

Location

Future

- Are there volcanoes in the site region and how old are they?
- Is the tectonic setting consistent with future volcanism?

from Valentine and Perry (2009)



Initial Scoping: Lava flows in the Harrat Al Shamm (Jordan)

PVHA

Background


Conceptual model

Rates

Location

Future

Are there volcanoes in the site region and how old are they? Is the tectonic setting consistent with future volcanism? (Figure by WesternGeco, 2018, with permission from JAEC)



Consider potential for specific volcanic products in characterizing sources. Develop a conceptual model of potential volcanic activity based on geologic record, analog volcanic systems:

Phenomena	exclusionary?	migitation?
Opening of new vents	Yes	No
Sector Collapse	Yes	No
Pyroclastic density currents	Yes	No
Lava flows	Yes	No
Lahar	Yes	Yes
Tephra fallout	No	Yes
Volcanic gases	No	Yes
Volcanic earthquakes	No	Yes

Characterize Sources: Volcanism in the Eifel volcanic field (Germany)

PVHA

Background

Conceptual model


Rates

Location

Future



Pyroclastic surges, tephra fallout, and damming of Rhine:



From Park and Schmincke, 1997

Conceptual Model: Post-collisional volcanism (Armenia)

PVHA


Background

Conceptual model

Rates

Location

Future



from L. J. Connor et al., 2012

Conceptual Model: volcanic hazards on Ischia (Italy)

PVHA

Background



Conceptual model

Rates

Location

Future

What volcanic hazards do residents of Ischia face given the nature of volcanic activity during the last 150 ka?



From Selva et al., 2019, JAV

Screening Hazards based on geologic record: Armenia

PVHA

Background


Conceptual
model

Rates

Location

Future

Given a volcanic eruption, is it possible specific volcanic phenomena can reach the site?



Low-aspect ratio ignimbrites and lava flows reach the Armenia Nuclear Power Plant site.

Screening Hazards based on simulations: Tonila (Mexico)

PVHA

Background

Conceptual model

Rates


Location

Future

Given a volcanic eruption, is it possible specific volcanic phenomena can reach the site?



Based on simulations, lava flows from the summit of Volcán de Colima of 0.1 km^3 might reach the Tonila vicinity



Screening Hazards based on simulations: Tonila (Mexico)

PVHA

Background


Conceptual model

Rates


Location

Future

Given a volcanic eruption, is it possible specific volcanic phenomena can reach the site?



energy cone model



tephra simulation

Screening Hazards: Ikata (Japan)

PVHA

Background

Conceptual
model

Rates

Location

Future

Kazuhiro Nakamoto, President, Japan Federation of Bar Associations

Today, the Hiroshima High Court handed down a temporary injunction compelling Shikoku Electric Power Corporation to stop operation of the No. 3 reactor of the Ikata Nuclear Power Plant. The decision was made according to the evaluation procedures in the volcanic eruption guidelines set by the Nuclear Regulation Authority (NRA). It was found that it was difficult to judge whether the volcanic activity of the Mt. Aso caldera, located 130 kilometers away from the Ikata NPP, was weak enough during the operation of the reactor. As it is impossible to estimate how big an eruption of Mt. Aso would be, the judgment took the largest past eruption of Mt. Aso “Aso-4” (about 90,000 years ago) (volcanic explosivity index 7) as the basis for its assumption. **The court found that it cannot conclude that the Aso-4 pyroclastic flow was very unlikely to reach Ikata NPP, and therefore judged that the Ikata NPP was not located in an appropriate location.**

PVHA


Background

Conceptual model

Rates

Location

Future



Models suggest 44 ± 7 yr for European tephra clouds

PVHA

Background


Conceptual model

Rates



Location

Future

When eruption frequency is stationary, it is possible to apply univariate statistical models to estimate recurrence rates



Swindles et al. (2011) *Geology*.
 Watson et al. (2017) *EPSL*.
 Swindles et al. (2017) *Geology*.





Models suggest 44 ± 7 yr for European tephra clouds

PVHA

Calculate stationarity within some confidence interval:

$$\hat{\lambda} = \frac{N - 1}{t_o - t_y}$$


$$CI_{95\%} = \hat{\lambda}t \pm 1.36/\sqrt{N}$$



Bebbington, 2013, Connor et al. 2015, Watson et al. ,2017

$$P[N \geq 1] = 1 - \exp[-\lambda \Delta t]$$

Frequency of explosive eruptions at Momotombo volcano (Nicaragua) measured during Feb–April, 2016



Probability model must (1) use a subset of data, or (2) detrend the data, or (3) use a cluster or renewal model). One cannot apply a univariate model, like an exponential model, to nonstationary distributions.

A Volcano Eruption Age Model for Mars

PVHA


Background

Conceptual model

Rates

Location

Future



Vents and lava flows in the caldera of Arsia Mons are among Mars' youngest volcanoes. How do we constrain the timing of these eruptions? *Richardson et al., EPSL, 2017*

Map relations among lava flows reveal stratigraphy

PVHA


Background

Conceptual model

Rates

Location

Future



Context Imager (CTX) datasets are used to map stratigraphic relationships within the caldera at the summit of Arsia Mons.

A directed graph of age and stratigraphic relationships

PVHA


Background

Conceptual model

Rates

Location

Future



Ages estimated (with high uncertainty!) from crater density

Randomly sample ages of all events using directed graph
($M = 10000$ times),

Volcano i of total N formed by event \hat{e}_i ,


For each set of age estimates, j , for N volcanoes, the cumulative distribution is:

$$X_j(T) = \sum_{i=1}^N P[\hat{e}_{i,j}, t < T]$$

where $P[\hat{e}_{i,j}, t < T] = 0$ if $T < \hat{e}_{i,j}$ and $P[\hat{e}_{i,j}, t < T] = 1$ if $T \geq \hat{e}_{i,j}$

$$E(X) = \frac{1}{M} \sum_{j=1}^M X_j(T)$$

$$R(X) = \frac{\Delta E(X)}{\Delta t}$$



Based on Monte Carlo simulation using age estimates and stratigraphic information

Age distribution of events is improved by using directed graph with Monte Carlo simulation

PVHA


Background

Conceptual model

Rates

Location

Future



March 20, 2017

Mars Volcano, Earth's Dinosaurs Went Extinct About the Same Time



Estimating recurrence rate from the rock record

PVHA

Background

Conceptual model

Rates

Location

Future



System	Stage	Geochronometry, Ma	Geological Unit	Age, Ma, K-Ar	Age, Ma, Ar-Ar	Stage	Description
Quaternary	Upper Pleistocene, O ₃	0.0117	1. Q3-Q4, a,d,e,p				Alluvia, diluvia, eluvia, proluvia deposits, pebble, sand, sandy-loam, loam, rubble.
	Middle Pleistocene, Q ₂	0.126	2. Q2, a,f,g,m				Glacial and fluvio-glacial deposits, moraines
		3. Q2, B-BA	0.45-0.53	0.614	IV	Basaltic trachyandesite and basaltic-andesite lava flows, Tonsakar Ashconic, Dambakar	
		4. Q2, TD		0.75		Trachydicite lava flows of Cahkisar volcano (Pohr Bogatu)	
		5. Q2, b,f,g,m				Glacial and fluvio-glacial deposits, moraines	
		6. Q2, TA-TD	0.73-0.54			Trachyandesites, trachydicites, of near the summit plateau	
		7. Q2, Ig,A				Tuffs (ignimbrites) of Artk horizon	
		8. Q2, B-BA				Basaltic and basaltic-andesite lava flows of Kakavasar, SW slopes in Aragats	
		9. Q2, B-BA				Basaltic and basaltic-andesite lava flows of Sharshar (Gogak) group of volcanoes (N. Aragats)	
		10. Q2, TB-TBA	0.74-0.90			trachybasaltic and basaltic trachyandesite lava flows of Uerzhay plateau, K-Ar	
		11. Q2, D,P,I,g		0.49		trachyandesite and andesite plateau, Dacic lava flows, Pirran eruption pumice fallout deposits, hydrocalcicite ignimbrites	
	12. Q2, Ig, YG-BS		0.65-0.66		ignimbrite tuffs of Yerevan-Gyumri type and Byurakan-Shamiram subtype		
	13. Q1-2, B-BA				Basalts, Basaltic-andesites covering Arvi type tuff		
	14. Q1-2, P-Ig				Pumice ignimbrite tuffs of Arvi type		
	0.781	15. Q1-2, TA-TD,P			Trachyandesites, trachydicites of slopes of Aragats, in South part covered by trind dacites		
	16. Q1-2, A-D				Pumice eruption fallout deposits of Parnashan		
	17. Q1-2, LS				Andesites, dacites of Duzan and Byurakan type		
	18a-b. Q1-2, TBA-TA	0.91-1.10	0.71-1.32	0.809	II	Upper unit of lake sediments deposits of Ararat and Shrak valleys and Apsaran depression	
	19. Q1-2, A-D	0.92-0.99	0.902		Basaltic-andesites, basaltic-trachyandesites, andesites and trachyandesites of Shamiram and Eghvard plateaus, Barmbarod, Grawko and other similar zones		
	20. Q1, TD-Rh	1.45-1.60			Basaltic trachyandesites of Sardapat structure		
	1.806	21. N ₁ ⁺ , Q1, B-BA			Trachydicites and rhyolites of Ararat volcano		
	22. N ₁ ⁺ , B-BA (D)	2.20-2.50			Basalts and basaltic-andesites of S and SW Aragats		
				Dolerite basalts and basaltic andesite			

Recurrence rate: Known eruptions by volume

PVHA


Background

Conceptual model

Rates


Location

Future



- TB-TA: trachy-basalt trachy-andesite
- TD: trachy-dacite

Main concern: is there no activity in the last 400 ka? Or is there a lack of preservation of smaller eruptions?



- Cumulative distribution function of Aragats eruptions.
- Steady-state activity until about 0.5 Ma, after which no eruptions are identified.

Recurrence rate: weighting alternative models with expert judgment

PVHA


Background

Conceptual model

Rates

Location

Future



Node	Recurrence Rate (yr^{-1})	Wt
$N_{3,1}$	$> 5 \times 10^{-5}$	0
$N_{3,2}$	$> 0.5 - 5 \times 10^{-5}$	0.1
$N_{3,3}$	$0.9 - 5 \times 10^{-6}$	0.45
$N_{3,4}$	$< 9 \times 10^{-7}$	0.45

logic tree for pdcs impacting the site

PVHA


Background

Conceptual model

Rates

Location

Future



PVHA

Background

Conceptual model

Rates

Location

Future

Range (High/Low)	Weight ($w_{2,1-4}$)	RR (yr^{-1})	Weight ($1 - w_{3,1-3}$)	Weight ($1 - w_4$)	Weighted Probability
H	0.10	5e-05	0.67	1	3.3×10^{-6}
L	0.10	5e-06	0.33	1	1.6×10^{-7}
H	0.45	5e-06	0.10	1	2.2×10^{-7}
L	0.45	9e-07	0.10	1	4×10^{-8}
H	0.45	9e-07	0.01	1	4×10^{-9}

$$\text{Aggregate Annual Probability} = 2 \times 10^{-7} - 3.5 \times 10^{-6}$$

Spatial density of volcanic vents

PVHA


Background

Conceptual
model

Rates

Location

Future



spatial density of volcanic vents

PVHA


Background

Conceptual
model

Rates

Location

Future



Probability of Opening of New Vents and Lava Inundation

PVHA


Background

Conceptual model

Rates

Location

Future



Hazard	Annual Probability (vents)	Annual Probability (Events)
Eruption on the ESRP	5.7×10^{-4}	2.6×10^{-4}
Eruption in INL	1.2×10^{-4}	6.2×10^{-5}
Lava Inundation of INL	1.8×10^{-4}	8.4×10^{-5}

see Gallant et al (2018)

PVHA

Background

Conceptual model

Rates

Location

Future

- Long-term volcanic hazard assessment must become more widespread to plan for volcanic activity, in all its forms, before it affects communities and infrastructure.
- Probabilistic volcanic hazard assessment relies on a simple hierarchical structure (e.g., logic trees). What new structures should emerge?
- PVHA places a premium on geologic data collection, especially radiometric age determinations and mapping, and numerical models
- Major challenge is to improve monitoring to identify potentially active volcanic systems before “unrest” .

- *IAEA*: S. Aramaki, W. Aspinall, S. Charbonnier, A. Chigama, O. Coman, L. J. Connor, A. Costa, L. Courtland, H. Delgado Granados, A. Godoy, B. Hill, C. Jaupart, J.-C. Komorowski, A. McBirney, S. McNutt, K. Meliksetian, S. Nakada, C. Newhall, G. Pasquare, I. Savov, S. Self, Y. Uchimyama, T. Wilson
- Jacob Richardson, Lis Gallant, Graeme Swindles, Elizabeth Watson, Armando Saballos, Kristy Murphy, Mike Sheridan,